

- The Clipboard will hold a maximum of 500 items.
- Clipboard items will be lost after one hour of inactivity.

Display	Summary	Sort	Save	Text	Clip Remove	Order
Show: 20		Items 1-12 of 12				One page

PubMed Services

Related Resources

5661084

5725351
5795564
5670149

□ 1: [Mackay F, Bourdon PR, Griffiths DA, Lawton P, Zafari M, Sizing ID, Miatkowski K, Ngam-ek A, Benjamin CD, Hession C, Ambrose CM, Meier W, Browning JL.](#) [Related Articles](#)
Cytotoxic activities of recombinant soluble murine lymphotoxin-alpha and lymphotoxin-alpha beta complexes.
J Immunol. 1997 Oct 1;159(7):3299-310.
PMID: 9317128 [PubMed - indexed for MEDLINE]

□ 2: [Mackay F, Majeau GR, Lawton P, Hochman PS, Browning JL.](#) [Related Articles](#)
Lymphotoxin but not tumor necrosis factor functions to maintain splenic architecture and humoral responsiveness in adult mice.
Eur J Immunol. 1997 Aug;27(8):2033-42.
PMID: 9295042 [PubMed - indexed for MEDLINE]

□ 3: [Browning JL, Dougas I, Ngam-ek A, Bourdon PR, Ehrenfels BN, Miatkowski K, Zafari M, Yampaglia AM, Lawton P, Meier W, et al.](#) [Related Articles](#)
Characterization of surface lymphotoxin forms. Use of specific monoclonal antibodies and soluble receptors.
J Immunol. 1995 Jan 1;154(1):33-46.
PMID: 7995952 [PubMed - indexed for MEDLINE]

□ 4: [Cosman D.](#) [Related Articles](#)
A family of ligands for the TNF receptor superfamily.
Stem Cells. 1994 Sep;12(5):440-55. Review.
PMID: 7528588 [PubMed - indexed for MEDLINE]

□ 5: [Armitage RJ.](#) [Related Articles](#)
Tumor necrosis factor receptor superfamily members and their ligands.
Curr Opin Immunol. 1994 Jun;6(3):407-13. Review.
PMID: 7917108 [PubMed - indexed for MEDLINE]

□ 6: [Crowe PD, VanArsdale TL, Walter BN, Ware CF, Hession C, Ehrenfels B, Browning JL, Din WS, Goodwin RG, Smith CA.](#) [Related Articles](#), [Nucleotide](#), [OMIM](#), [Protein](#)
A lymphotoxin-beta-specific receptor.
Science. 1994 Apr 29;264(5159):707-10.
PMID: 8171323 [PubMed - indexed for MEDLINE]

□ 7: [Browning JL, Ngam-ek A, Lawton P, DeMarinis J, Tizard R, Chow EP, Hession C, O'Brine-Greco B, Foley SF, Ware CF.](#) [Related Articles](#), [Nucleotide](#), [OMIM](#), [Protein](#)
Lymphotoxin beta, a novel member of the TNF family that forms a heteromeric complex with lymphotoxin on the cell surface.

Cell. 1993 Mar 26;72(6):847-56.
PMID: 7916655 [PubMed - indexed for MEDLINE]

8: [Hochman PS, Majeau GR, Mackay F, Browning JL.](#)

Related Articles

Proinflammatory responses are efficiently induced by homotrimeric but not heterotrimeric lymphotoxin ligands.

J Inflamm. 1995-96;46(4):220-34.

PMID: 8878796 [PubMed - indexed for MEDLINE]

9: [Gerspach J, Gotz A, Zimmermann G, Kolle C, Bottinger H, Grell M.](#)

Related Articles

Detection of membrane-bound tumor necrosis factor (TNF): an analysis of TNF-specific reagents.

Microsc Res Tech. 2000 Aug 1;50(3):243-50.

PMID: 10891890 [PubMed - indexed for MEDLINE]

10: [Ettinger R, Browning JL, Michie SA, van Ewijk W, McDevitt HO.](#)

Free in PMC, Related Articles

Disrupted splenic architecture, but normal lymph node development in mice expressing a soluble lymphotoxin-beta receptor-IgG1 fusion protein.

Proc Natl Acad Sci U S A. 1996 Nov 12;93(23):13102-7.

PMID: 8917551 [PubMed - indexed for MEDLINE]

11: [Gruss HJ, Dower SK.](#)

Related Articles

The TNF ligand superfamily and its relevance for human diseases.

Cytokines Mol Ther. 1995 Jun;1(2):75-105. Review.

PMID: 9384666 [PubMed - indexed for MEDLINE]

12: [Neumann B, Luz A, Pfeffer K, Holzmann B.](#)

Related Articles

Defective Peyer's patch organogenesis in mice lacking the 55-kD receptor for tumor necrosis factor.

J Exp Med. 1996 Jul 1;184(1):259-64.

PMID: 8691140 [PubMed - indexed for MEDLINE]

Display	Summary	Sort	Save	Text	Clip	Remove	Order
Show:	20	▼	Items 1-12 of 12				One page

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

sparc-sun-solaris2.8 Oct 4 2001 12:24:52